RS = Disease Response and Clin Classification

RSCAT = BPRS

BRIEF PSYCHIATRIC RATING SCALE (BPRS)

D NI				,	m 1 . D .	RSDTC
Patient Name C/RSSTRESN]				Today's Date	
Please enter tr	e score fo	r the term that	best describes th	e patient's condition. RSOR	RES	
		Not present,	2 = Very mild,	3 = Mild, 4 = Moderate, 5 = 1	Moderately severe,	6 = Severe,
7 = Extremely	/ severe					
Score						
1.		FIC CONCE pation with ph		r of physical illness, hypochondi	riasis. RSTESTC	D = BPR0201
2.	ANXIE Worry, f		ern for present or	future, uneasiness.	RSTESTC	D = BPR0202
3.		ONAL WIT		on deficiency in relating to othe		D = BPR0203
4.			SORGANIZAT afused, disconnec	ION ted, disorganized, disrupted.	RSTESTC	D = BPR0204
5.		FEELINGS ne, shame, ren	norse for past beh	avior.	RSTESTO	D = BPR0205
6.	TENSION Physical		nifestations of n	ervousness, over-activation.	RSTEST	CD = BPR020
7.			D POSTURIN tural motor behav	G vior (not including tic).	RSTEST	CD = BPR020
8.		DIOSITY ated self-opinio	on, arrogance, co	nviction of unusual power or abi	ilities. RSTEST	CD = BPR020
9.		SSIVE MOO sadness, despo	D ndency, pessimisi	n.	RSTEST	CD = BPR020
10.	HOSTI Animos		belligerence, disc	lain for others.	RSTEST	CD = BPR02
11.		CIOUSNESS , belief others	harbor malicious	or discriminatory intent.	RSTES1	CD = BPR02
12.			BEHAVIOR ormal external st	imulus correspondence.	RSTEST	CD = BPR02
13.		R RETARDA weakened mo		h, reduced body tone.	RSTEST	CD = BPR02
14.		OPERATIVE ce, guardednes	ENESS ss, rejection of au	thority.	RSTES1	CD = BPR02
15.			GHT CONTEN		RSTES	TCD = BPR02
16.		FED AFFECT l emotional to		formal intensity of feelings, flatne	ess.	TCD = BPR02
17.	EXCIT: Heighte		tone, agitation,	increased reactivity.	RSTES	TCD = BPR02
18.		IENTATION		for person, place or time.	RSTES	TCD = BPR0

BRIEF PSYCHIATRIC RATING SCALE (BPRS)

Instructions for the Clinician:

The Brief Psychiatric Rating Scale (BPRS) is a widely used instrument for assessing the positive, negative, and affective symptoms of individuals who have psychotic disorders, especially schizophrenia. It has proven particularly valuable for documenting the efficacy of treatment in patients who have moderate to severe disease.

It should be administered by a clinician who is knowledgeable concerning psychotic disorders and able to interpret the constructs used in the assessment. Also considered is the individual's behavior over the previous 2-3 days and this can be reported by the patient's family.

The BPRS consists of 18 symptom constructs and takes 20-30 minutes for the interview and scoring. The rater should enter a number ranging from 1 (not present) to 7 (extremely severe). 0 is entered if the item is not assessed.

First published in 1962 as a 16-construct tool by Drs. John Overall and Donald Gorham, the developers added two additional items, resulting in the 18-item scale used widely today to assess the effectiveness of treatment.

BPRS Scoring Instructions: R

RSTESTCD = BPR0219

Sum the scores from the 18 items. Record the total score and compare the total score from one evaluation to the next as the measure of response to treatment.