

RS=Disease Response and Clin Classification

RSCAT=KDIGO AKI

KDIGO Staging For Acute Kidney Injury

RSSTRESC/RSSTRESN

Instructions	Item	Answer
<p>Check the correct box to indicate the patient's Acute Kidney Injury stage.</p>	<p>KDIGO Stage</p> <p>RSTESTCD=KDIGO101</p>	<p>RSORRES</p> <p><input type="checkbox"/> Stage 1</p> <p>Serum creatinine criteria: 1.5-1.9 times baseline OR ≥ 0.3 mg/dL (≥ 26.5 $\mu\text{mol/L}$) increase Urine output criteria: < 0.5 mL/kg/h for 6-12 hours</p> <p><input type="checkbox"/> Stage 2</p> <p>Serum creatinine criteria: 2.0-2.9 times baseline Urine output criteria: < 0.5 mL/kg/h for ≥ 12 hours</p> <p><input type="checkbox"/> Stage 3</p> <p>Serum creatinine criteria: 3.0 times baseline OR Increase in serum creatinine to ≥ 4.0 mg/dL (≥ 353.6 $\mu\text{mol/L}$) OR Initiation of renal replacement therapy OR in patients < 18 years, decrease in eGFR to < 35 mL/min per 1.73 m^2 Urine output criteria: < 0.3 mL/kg/h for ≥ 24 hours OR Anuria for ≥ 12 hours</p>

Reference: : Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO Clinical Practice Guideline for Acute Kidney Injury. *Kidney inter., Suppl.* 2012;2: 1–138.

CDISC created this case report form to represent the KDIGO Stage for Acute Kidney Injury, which is in the public domain. This is not a validated CRF or an endorsement of the KDIGO. CDISC specifies how to structure the data that has been collected in a database, not what should be collected or how to conduct clinical assessments or protocols.